

S2 Table. Pathological findings of subsolid lung adenocarcinomas

Variable	A(-) I(-) (n=120)	A(+) I(-) (n=26)	A(-) I(+) (n=117)	A(+) I(+) (n=141)	p-value
Pathological T category					
T1	118 (98.3)	24 (92.3)	94 (80.3)	96 (68.1)	< 0.001
T2	2 (3.4)	2 (7.8)	22 (18.8)	42 (29.8)	
T3-4	0 (0.0)	0 (0.0)	1 (0.9)	3 (2.1)	
Pathological N category					
Nx	28 (23.3)	5 (19.2)	17 (14.5)	23 (16.3)	0.235
N0	90 (75.0)	21 (80.8)	96 (82.1)	106 (75.2)	
N1	1 (0.8)	0 (0.0)	1 (0.9)	7 (5.0)	
N2	1 (0.8)	0 (0.0)	3 (2.6)	5 (3.6)	
Overall pathological category					
I	118 (98.3)	25 (96.2)	108 (92.3)	123 (87.2)	0.038
II	1 (0.8)	1 (3.9)	6 (5.1)	12 (8.5)	
III	1 (0.8)	0 (0.0)	3 (2.6)	6 (4.3)	
Vascular invasion	0 (0.0)	0 (0.0)	17 (14.5)	20 (14.2)	< 0.001
Lymphatic invasion	5 (4.2)	2 (7.7)	25 (21.4)	31 (22.0)	< 0.001

Values are presented as number (%). A, attachment; I, indentation.