

S2 Table. Geometric mean titers (GMTs) of opsonophagocytic activity (OPA) stratified by chemotherapeutic regimen

Serotype	Arm	Pre-vaccination			Post-vaccination		
		GMT	95% CI	p-value	GMT	95% CI	p-value
Fluoropyrimidine (n=25)							
5	A	4	2-8	0.490	574	170-1,942	0.303
	B	6	3-15		164	16-1,724	
6B	A	496	90-2,728	0.956	13,489	9,658-18,838	0.048
	B	530	71-3,983		4,693	1,697-12,976	
18C	A	163	54-493	0.718	7,059	4,211-11,833	0.239
	B	112	15-843		2,267	305-16,839	
19A	A	716	231-2,221	0.380	11,266	7,335-17,303	0.015
	B	344	83-1,430		2,658	925-7,633	
Fluoropyrimidine+Oxaliplatin (n=62)							
5	A	5	3-8	0.689	247	98-622	0.183
	B	4	3-7		510	265-982	
6B	A	537	145-1,992	0.920	6,972	3,823-12,714	0.030
	B	498	207-1,201		12,910	8,501-14,391	
18C	A	75	32-177	0.583	3,181	1,655-6,115	0.104
	B	102	48-217		6,057	3,726-9,847	
19A	A	325	129-820	0.881	6,001	2,830-12,724	0.284
	B	300	157-573		8,896	6,527-12,123	

Pre- and post-vaccination GMTs with 95% confidence intervals (CIs) in the two study arms, stratified by chemotherapeutic regimen (fluoropyrimidine vs. fluoropyrimidine+oxaliplatin).