

S3 Table. Univariate and multivariate analyses for overall survival (waiting time as categorical variable based on median of 14 days)

	Univariate analysis			Multivariate analysis		
	HR	95% CI	p-value	HR	95% CI	p-value
Interval (day)						
0-14	1			1		
15-60	1.405	1.009-1.955	0.044	1.513	1.073-2.134	0.018
Age (yr)	1.015	0.999-1.030	0.059	1.015	0.999-1.031	0.066
FIGO stage			0.001			0.001
I	1			1		
II	1.102	0.528-2.304	0.796	1.138	0.540-2.399	0.734
III	2.015	0.941-4.316	0.071	2.273	1.056-4.893	0.036
IV	2.255	1.040-4.890	0.039	2.386	1.097-5.187	0.028
Cell type			< 0.001			< 0.001
SCC	1			1		
AD/ADS	2.337	1.540-3.548	< 0.001	2.224	1.460-3.388	< 0.001
Others	4.550	1.440-14.372	0.010	5.067	1.548-16.589	0.007
Chemotherapy ^{a)}						
Single agent	1			1		
Combination	1.279	0.917-1.784	0.147	1.267	0.883-1.820	0.200
Early discontinuation ^{b)}						
Yes	1			1		
No	0.427	0.274-0.666	<0.001	0.436	0.277-0.686	<0.001

AD, adenocarcinoma; ADS, adenosquamous cell carcinoma; Others, small cell, clear cell, poorly differentiated carcinoma; SCC, squamous cell carcinoma. ^{a)}Chemotherapy during CCRT, ^{b)}Early discontinuation of chemotherapy during CCRT (e.g., 1 or 2 cycles for combination chemotherapy or 1, 2, or 3 for single agent).