

S3 Table. Patient characteristics according to adjuvant chemotherapy status

	No adjuvant chemotherapy (n=43)	Adjuvant chemotherapy (n=170)	p-value
Age (yr)	69.77±7.29	64.17±8.99	< 0.001
Male sex	35 (81.4)	148 (87.6)	0.321
Clinical T category			0.518
T1/T2/T3/T4	24 (55.8)/18 (41.9)/0/1 (2.3)	86 (50.6)/73 (42.9)/8 (4.78)/3 (1.8)	
Clinical N category			0.068
N0/N1/N2/N3	36 (83.7)/3 (7.0)/3 (7.0)/1 (2.3)	124 (72.9)/29 (17.1)/17 (10.0)/0	
Pathologic T category			0.422
T1/T2/T3/T4	22 (51.2)/16 (37.2)/5 (11.6)/0	67 (39.4)/79 (46.5)/20 (11.8)/4 (2.4)	
Pathologic N stage			0.032
Nx/N0/N1/N2/N3	1 (2.3)/32 (74.5)/5 (11.6)/4 (9.3)/1 (2.3)	4 (2.3)/93 (54.8)/41 (24.1)/32 (18.8)/0	
Extent of resection			0.850
Wedge resection	7 (16.3)	19 (11.2)	
Segmentectomy	3 (7.0)	11 (6.5)	
Lobectomy or bilobectomy	31 (72.1)	127 (74.7)	
Pneumonectomy	2 (4.7)	13 (7.6)	
Mediastinal lymph node dissection			0.188
Systematic MLND	38 (88.4)	160 (94.1)	
Less than systematic MLND	5 (11.6)	10 (5.9)	

Values are presented as mean±standard deviation or number (%). MLND, mediastinal lymph node dissection.