

**S3 Table.** Adverse events occurring in at least 10% of patients in either group of patients with CRC (n=366)

	Female (n=178, 48.6%)		Male (n=188, 51.4%)		p-value	
	Any grade	Grade $\geq$ 3	Any grade	Grade $\geq$ 3	Any grade	Grade $\geq$ 3
Neutropenia	135 (75.8)	47 (26.4)	115 (61.2)	42 (22.3)	0.003	0.365
Anemia	115 (64.6)	2 (1.1)	114 (60.6)	5 (2.7)	0.451	0.450
Peripheral neuropathy	99 (55.6)	2 (1.1)	94 (50.0)	0	0.282	0.236
Thrombocytopenia	97 (54.5)	4 (2.2)	102 (54.3)	3 (1.6)	0.963	0.717
AST elevation	71 (39.9)	0	46 (24.5)	2 (1.1)	0.002	0.499
Anorexia	69 (38.8)	1 (0.6)	68 (36.2)	2 (1.1)	0.608	> 0.99
Nausea	61 (34.3)	25 (14.0)	38 (20.2)	16 (8.5)	0.002	0.093
Hand-foot syndrome	57 (32.0)	1 (0.6)	62 (33.0)	0	0.845	0.486
ALT elevation	56 (31.5)	2 (1.1)	44 (23.4)	4 (2.1)	0.084	0.686
Diarrhea	37 (20.8)	1 (0.6)	37 (19.7)	2 (1.1)	0.792	> 0.99
Abdominal pain	38 (21.3)	3 (1.7)	24 (12.8)	1 (0.5)	0.029	0.360
Stomatitis	37 (20.8)	0	22 (11.7)	1 (0.5)	0.018	> 0.99
Fatigue	34 (19.1)	0	34 (18.1)	0	0.803	-
Vomiting	32 (18.0)	20 (11.2)	15 (8.0)	8 (4.3)	0.004	0.012
Skin rash	25 (14.0)	18 (9.6)	1 (0.5)	0	0.184	> 0.99
Constipation	18 (0.1)	0	13 (6.9)	0	0.272	-

Values are presented as number (%). ALT, alanine aminotransferase; AST, aspartate aminotransferase.