

S6 Table. Adverse events occurring in at least 10% of patients in either group of patients who received platinum-containing regimen and had unscheduled outpatient clinic visits (n=95)

	Female (n=52)		Male (n=43)		p-value	
	Any grade	Grade \geq 3	Any grade	Grade \geq 3	Any grade	Grade \geq 3
Neutropenia	33 (34.7)	8 (8.4)	25 (26.3)	7 (7.4)	0.596	0.905
Anemia	34 (35.8)	1 (1.1)	30 (31.6)	0 (0.0)	0.650	-
Peripheral neuropathy	28 (29.5)	1 (1.1)	27 (28.4)	1 (1.1)	0.379	> 0.99
Anorexia	22 (23.2)	1 (1.1)	14 (14.7)	0	0.330	> 0.99
Thrombocytopenia	20 (21.1)	1 (1.1)	21 (22.1)	2 (2.1)	0.309	0.588
AST elevation	17 (17.9)	0	13 (13.7)	0	0.797	-
Hand-foot syndrome	16 (16.8)	0	18 (18.9)	0	0.262	-
Nausea	15 (15.8)	2 (2.1)	7 (7.4)	1 (1.1)	0.148	> 0.99
Stomatitis	15 (15.8)	-	2 (2.1)	-	0.002	-
Abdominal pain	12 (12.6)	3 (3.2)	4 (4.2)	0	0.074	0.249
ALT elevation	11 (11.6)	1 (1.1)	8 (8.4)	1 (1.1)	0.757	> 0.99
Vomiting	11 (11.6)	3 (3.2)	3 (3.2)	1 (1.1)	0.052	0.405
Fatigue	11 (11.6)	0	11 (11.6)	0	0.611	—

Values are presented as number (%). ALT, alanine aminotransferase; AST, aspartate aminotransferase.