

S11 Table. Univariate and multivariate analyses of risk factors associated with recurrence (n=203) in *EGFR*-mutant stage IB NSCLC patients after curative surgery

Parameter	Univariate HR (95% CI)	p-value	Multivariate HR (95% CI)	p-value
Sex				
Female vs. Male (Ref)	0.52 (0.22-1.22)	0.131	-	-
Age (yr)				
> 65 vs. ≤ 65 (Ref)	1.13 (0.56-2.27)	0.736	-	-
BMI				
Per 1 unit increase	1.05 (0.93-1.19)	0.417	-	-
Smoking				
Ever vs. never (Ref)	0.52 (0.20-1.35)	0.178	-	-
DLco (per)				
Per 1 percentage increase	0.99 (0.97-1.01)	0.191	0.98 (0.96-1.00)	0.078
Pleural invasion				
Visceral pleural invasion vs. no pleural invasion (Ref)	2.68 (1.27-5.64)	0.010	2.75 (1.31-5.75)	0.007
Surgery type				
Lobectomy vs. wedge resection	0.39 (0.08-1.91)	0.248	-	-

BMI, body mass index; CI, confidence interval; DLco, diffusing capacity of the lungs for carbon monoxide; EGFR, epidermal growth factor receptor; HR, hazard ratio; NSCLC, non-small cell lung cancer.