

S12 Table. Univariate and multivariate logistic regression analyses for early recurrence in *EGFR*-mutated NSCLC patients after curative surgery (n=867)

Parameter	Univariate OR (95% CI)	p-value	Multivariate OR (95% CI)	p-value
Sex				
Female vs. male (Ref)	1.42 (0.66-3.04)	0.370	-	-
Age (yr)				
> 65 vs. ≤ 65 (Ref)	1.40 (0.72-2.69)	0.320	-	-
BMI				
Per 1 unit increase	0.94 (0.86-1.04)	0.223	-	-
Smoking				
Ever vs. never (Ref)	2.85 (1.33-6.12)	0.007	2.24 (1.18-4.23)	0.013
DLco (per)				
Per 1 percentage increase	1.0 (0.98-1.01)	0.588		
pT				
T3 vs. T1 (Ref)	4.42 (1.39-14.08)	0.012	5.11 (1.75-14.93)	0.003
pN				
N2 vs. N0 (Ref)	3.14 (1.42-6.96)	0.005	3.11 (1.44-6.68)	0.004
Pleural invasion				
Visceral pleural invasion vs. no pleural invasion (Ref)	1.43 (0.68-3.03)	0.346	-	-

BMI, body mass index; CI, confidence interval; DLco, diffusing capacity of the lungs for carbon monoxide; EGFR, epidermal growth factor receptor; NSCLC, non-small cell lung cancer; OR, odds ratio.