

**S9 Table.** Univariate and multivariate analyses of risk factors associated with recurrence (n=390) in *EGFR*-mutant stage IA NSCLC patients after curative surgery

<b>Parameter</b>	<b>Univariate HR (95% CI)</b>	<b>p-value</b>	<b>Multivariate HR (95% CI)</b>	<b>p-value</b>
<b>Sex</b>				
Female vs. male (Ref)	0.48 (0.26-0.87)	0.017	0.46 (0.22-0.97)	0.043
<b>Age (yr)</b>				
> 65 vs. ≤ 65 (Ref)	1.40 (0.76-2.56)	0.283	-	-
<b>BMI</b>				
Per 1 unit increase	0.98 (0.89-1.08)	0.681	-	-
<b>Smoking</b>				
Ever vs. never (Ref)	1.49 (0.78-2.87)	0.230	-	-
<b>DLco (per)</b>				
Per 1 percentage increase	0.99 (0.97-1.01)	0.156	-	-
<b>pT</b>				
T1b vs. T1a (Ref)	1.07 (0.58-1.98)	0.832	-	-
<b>Surgery type</b>				
Lobectomy vs. wedge resection	0.40 (0.19-0.85)	0.018	0.35 (0.14-0.90)	0.029

BMI, body mass index; CI, confidence interval; DLco, diffusing capacity of the lungs for carbon monoxide; EGFR, epidermal growth factor receptor; HR, hazard ratio; NSCLC, non-small cell lung cancer.